

JOURNAL OF ALGEBRA 92, 283–302 (1985)

The Fundamental Representation of a Strongly Regular Baer Semigroup

K. S. S. NAMBOORIPAD

*Department of Mathematics, University of Kerala,
Kariavattom 695 581, India*

AND

F. J. PASTIJN

*Dienst Hogere Meetkunde, Rijksuniversiteit te Gent,
Krijgslaan 281, B-9000 Gent, Belgium*

Communicated by G. B. Preston

Received February 7, 1983

A bounded partially ordered set I is a complemented modular lattice if and only if the semigroup $B(I)$ of all strongly range-closed residuated transformations of I is a regular semigroup coordinatizing I . If S is any strongly regular Baer semigroup coordinatizing I , then the Janowitz representation of S maps S homomorphically onto a full regular subsemigroup of $B(I)$. It is shown that the Janowitz representation of S is equivalent to Hall's (or Grillet's) fundamental representation.

© 1985 Academic Press, Inc.

In this paper we use the notation and the terminology of [2, 4, 5, 14]. We begin by recalling some definitions of [5] (see also [13]) regarding cross-connections.

If P is a partially ordered set and $x \in P$, then $P(x)$ denotes the principal order ideal of P generated by x . Let P and Q be partially ordered sets. An order-preserving mapping $f: P \rightarrow Q$ is said to be *normal* if $\text{im } f = Q(a)$ for some $a \in Q$, and for every $x \in P$ there exists a $z \leq x$ such that $f|P(z)$ is an isomorphism of $P(z)$ onto $Q(xf)$. In particular, if f is normal, then there exists at least one element $b \in P$ such that f is an isomorphism of $P(b)$ onto $Q(a) = \text{im } f$. We denote by $M(f)$ the set of all elements $b \in P$ with this property.

If P , Q , and R are partially ordered sets, and if $f: P \rightarrow Q$ and $g: Q \rightarrow R$ are normal mappings, then fg is a normal mapping. Consequently, the set $S(P)$ of all normal transformations on a partially ordered set P is a semigroup under composition. Elements of $S(P)$ will be written as right operators, and

the elements of its left-right dual $S^{\text{op}}(P)$ will be written as left operators. Idempotents of $S(P)$ are called *normal retractions*. A principal ideal $P(a)$ of P is called a *normal retract* if $P(a) = \text{im } e$ for some normal retraction $e \in S(P)$. If every principal ideal of P is a normal retract, then P is called a *regular partially ordered set* [4, 5, 13].

An equivalence relation ρ on a poset P is said to be *normal* if there exists a normal mapping $f \in S(P)$ such that $\ker f = f f^{-1} = \rho$. The poset (under the reverse of inclusion) of all normal equivalences on P will be denoted by P° . Recall from [5] that when P is regular, then so is P° .

If f and g are normal mappings, with $\text{dom } f = \text{dom } g = P$, $\ker f = \ker g$, then it is easy to see that $M(f) = M(g)$. So with each $\rho \in P^\circ$ we may associate the subset $M(\rho)$ defined by $M(\rho) = M(f)$, where f is any normal mapping with $\ker f = \rho$. Remark that $a \in M(\rho)$ if and only if $P(a)$ intersects every ρ -class in exactly one element. If this is the case, then $P(a) \cap \rho(x)$ contains a single element which is minimal in its ρ -class, and the mapping $\varepsilon_\rho(\rho, a)$ which sends each $x \in P$ to the unique element in $P(a) \cap \rho(x)$ is a normal retraction with $\ker \varepsilon_\rho(\rho, a) = \rho$ and $\text{im } \varepsilon_\rho(\rho, a) = P(a)$. $\varepsilon_\rho(\rho, a)$ is called the *projection along ρ upon $P(a)$* .

If f is a normal mapping, with $\text{dom } f = P$, and $\ker f = \rho$, then for any $a \in M(f) = M(\rho)$, one can write $f = \varepsilon_\rho(\rho, a)\alpha$, where $\alpha = f|_{P(a)}$ is an isomorphism of $P(a)$ onto $\text{im } f$. This is called a *normal factorization* of f . Since for every normal mapping f we have $M(f) \neq \emptyset$, every normal mapping has at least one normal factorization, and there is a bijection between the set $M(f)$ and the set of normal factorizations of f .

A regular semigroup S is said to be *fundamental* if the identity congruence is the only idempotent-separating congruence on S . If P is a regular partially ordered set, then $S(P)$ is a fundamental regular semigroup such that $S(P)/\mathcal{L} \approx P$ and $S(P)/\mathcal{R} \approx P^\circ = \{\ker f \mid f \in S(P)\}$. Thus, $S(P)$ is a regular semigroup that coordinatizes P [5, 16].

PROPOSITION 1. *Let I and Λ be regular partially ordered sets and let $f: I \rightarrow \Lambda$ be a normal mapping. For $\sigma \in \Lambda^\circ$, define*

$$f^\circ(\sigma) = \ker(f\varepsilon_\Lambda(\sigma, u)) = \sigma f^{-1} \quad (1)$$

where $u \in M(\sigma)$. Then $f^\circ: \Lambda^\circ \rightarrow I^\circ$ is a normal mapping such that

$$\begin{aligned} \text{im } f^\circ &= I^\circ(\ker f) \\ M(f^\circ) &= \{\rho \in \Lambda^\circ \mid b \in M(\rho)\}, \quad \text{where } \text{im } f = \Lambda(b). \end{aligned} \quad (2)$$

If P , Q and R are regular partially ordered sets, and if $f: P \rightarrow Q$, and $g: Q \rightarrow R$ are normal mappings, then

$$(fg)^\circ = f^\circ g^\circ.$$

Proof. Let I and A be regular partially ordered sets, and suppose that $f: I \rightarrow A$ is a normal mapping. The mapping $f^\circ: A^\circ \rightarrow I^\circ$ which is given by (1) is order-preserving, and clearly $\text{im } f^\circ \subseteq I^\circ(\ker f)$.

If $\text{im } f = A(b)$, then we can choose $\rho \in A^\circ$ such that $b \in M(\rho)$, since A is a regular partially ordered set. We can put $f = f\epsilon_\Lambda(\rho, b)$. Also, if $a \in M(\ker f)$, then $f = \epsilon_I(\ker f, a)\alpha$, where $\alpha = f|I(a)$ maps $I(a)$ isomorphically onto $A(b)$.

Let $\sigma \in A^\circ$ and $c \in M(\sigma)$. Since $\epsilon_\Lambda(\rho, b)$ and $\epsilon_\Lambda(\sigma, c)$ are idempotents of the regular semigroup $S(A)$, the sandwich set $S(\epsilon_\Lambda(\rho, b), \epsilon_\Lambda(\sigma, c))$ is nonempty. Let $\epsilon_\Lambda(\sigma', b')$ be any element of this sandwich set. Then $\sigma' \supseteq \sigma$ and $b' \leq b$. We put $b'\alpha^{-1} = a' \in I(a)$. If $\tau = f^\circ(\sigma)$, we have

$$\begin{aligned}(u, v) \in \tau &\Leftrightarrow (uf) \epsilon_\Lambda(\sigma, c) = (vf) \epsilon_\Lambda(\sigma, c) \\ &\Leftrightarrow (uf) \epsilon_\Lambda(\sigma', b') = (vf) \epsilon_\Lambda(\sigma', b') \\ &\Leftrightarrow (u, v) \in f^\circ(\sigma'),\end{aligned}$$

and thus $\tau = f^\circ(\sigma')$. Let $\sigma'' \supseteq \sigma'$. Then there exists $b'' \in M(\sigma'')$ such that $b'' \leq b' \leq b$. Since $f^\circ(\sigma'') = \ker(f\epsilon_\Lambda(\sigma'', b''))$, $f\epsilon_\Lambda(\sigma'', b'')\alpha^{-1}$ is a normal retraction of I such that $\text{im}(f\epsilon_\Lambda(\sigma'', b'')\alpha^{-1}) = I(b''\alpha^{-1})$ and $\ker(f\epsilon_\Lambda(\sigma'', b'')\alpha^{-1}) = f^\circ(\sigma'')$. Thus for all $\sigma'' \supseteq \sigma'$ and $b'' \in M(\sigma'')$, with $b'' \leq b'$, one has

$$f\epsilon_\Lambda(\sigma'', b'') = \epsilon_I(f^\circ(\sigma''), b''\alpha^{-1})\alpha.$$

In particular we obtain

$$f\epsilon_\Lambda(\sigma', b') = \epsilon_I(f^\circ(\sigma'), a')\alpha = \epsilon_I(\tau, a')\alpha.$$

We now show that f° is an isomorphism of $A^\circ(\sigma')$ onto $I^\circ(\tau)$. Let $\tau'' \supseteq \tau$, and let $\sigma'' = \ker h$, where

$$h = \epsilon_\Lambda(\sigma', b')\alpha^{-1}\epsilon_I(\tau'', a''),$$

with

$$a'' \in M(\tau'') \text{ and } a'' \leq a' = b'\alpha^{-1} \leq a.$$

Then $\sigma'' \in A^\circ$ and $(x, y) \in f^\circ(\sigma'')$ if and only if $xfh = yfh$. But

$$\begin{aligned}xfh &= xf\epsilon_\Lambda(\sigma', b')\alpha^{-1}\epsilon_I(\tau'', a'') \\ &= x\epsilon_I(\tau, a')\alpha\alpha^{-1}\epsilon_I(\tau'', a'') \\ &= x\epsilon_I(\tau, a')\epsilon_I(\tau'', a'') \\ &= x\epsilon_I(\tau'', a'').\end{aligned}$$

Thus $xfh = yfh$ if and only if $(x, y) \in \tau''$, in other words, $f^\circ(\sigma'') = \tau''$. This proves that f° maps $A^\circ(\sigma')$ onto $I^\circ(\tau)$.

It is clear that $f^\circ|A^\circ(\sigma')$ preserves order. Assume that $\sigma_i \supseteq \sigma'$, $f^\circ(\sigma_i) = \tau_i \supseteq \tau$, $i = 1, 2$, and $\tau_1 \supseteq \tau_2$. Then $(u, v) \in \sigma_i$ if and only if $(u\epsilon_\Lambda(\sigma', b'), v\epsilon_\Lambda(\sigma', b')) \in \sigma_i$, $i = 1, 2$. Thus, putting $x = u\epsilon_\Lambda(\sigma', b')\alpha^{-1}$, $y = v\epsilon_\Lambda(\sigma', b')\alpha^{-1}$, then

$$\begin{aligned}(u, v) \in \sigma_2 &\Rightarrow (xf, yf) \in \sigma_2 \\ &\Rightarrow (x, y) \in f^\circ(\sigma_2) = \tau_2 \\ &\Rightarrow (x, y) \in f^\circ(\sigma_1) = \tau_1 \\ &\Rightarrow (xf, yf) \in \sigma_1 \\ &\Rightarrow (u, v) \in \sigma_1,\end{aligned}$$

from which $\sigma_2 \subseteq \sigma_1$. This proves that $f^\circ|A^\circ(\sigma')$ is an order isomorphism onto $I^\circ(\tau)$.

If in the above reasoning we put $\sigma = \rho$, then we obtain $\sigma' = \rho$ and $\tau = \ker f$. From this we infer $\text{im } f^\circ = I^\circ(\ker f)$, and we proved that f° is normal. From the above also follows that $M(f^\circ) \supseteq \{\rho \in A^\circ \mid b \in M(\rho)\}$. Let us now suppose that $\sigma \in M(f^\circ)$, i.e., $f^\circ|A^\circ(\sigma)$ is an isomorphism of $A^\circ(\sigma)$ onto $I^\circ(\ker f)$. The above reasoning shows that there exists a $\sigma' \supseteq \sigma$, $b' \in M(\sigma')$, $b' \leq b$, such that $f^\circ|A^\circ(\sigma')$ is an isomorphism of $A^\circ(\sigma')$ onto $I^\circ(\ker f)$. From this we have $\sigma = \sigma'$, whence $M(\sigma)$ contains an element b' , with $b' \leq b$. If $b' \neq b$, then $(b'\alpha^{-1}, b\alpha^{-1}) \in f^\circ(\sigma)$ and $(b'\alpha^{-1}, b\alpha^{-1}) \notin \ker f$. This is of course impossible since $f^\circ(\sigma) = \ker f$. Thus $b \in M(\sigma)$, and (2) follows.

We use the result proved above to reformulate the definition of cross-connections [5, 13].

Suppose that I and A are regular partially ordered sets and $\Gamma: A \rightarrow I^\circ$, $\Delta: I \rightarrow A^\circ$ are order preserving mappings. We say that $(f, g) \in S^{\text{op}}(I) \times S(A)$ is compatible with (Γ, Δ) if the following conditions hold:

- (C1) $\text{im } f = I(x)$, $\text{im } g = A(y) \Rightarrow \ker f = \Gamma(y)$, $\ker g = \Delta(x)$,
 (C2) the following diagrams commute:

$$\begin{array}{ccc} I & \xrightarrow{\Delta} & A^\circ \\ f \uparrow & & \uparrow g^\circ \\ I & \xrightarrow{\Delta} & A^\circ \end{array} \quad \begin{array}{ccc} A & \xrightarrow{\Gamma} & I^\circ \\ g \uparrow & & \uparrow f^\circ \\ A & \xrightarrow{\Gamma} & I^\circ \end{array}$$

(D1) (D2)

We observe that if I and A have identity elements, and if Γ and Δ preserve

identity elements, then (C2) implies (C1). In this case the identity relations on I and A are the identity elements of I° and A° , respectively.

In view of Proposition 1, the definition of cross-connections (see [5, 13, 16]) is equivalent to the following.

THEOREM 2. *Let I, A be regular partially ordered sets and let $\Gamma: A \rightarrow I^\circ$, $\Delta: I \rightarrow A^\circ$ be order preserving mappings. Then $[I, A; \Gamma, \Delta]$ is a cross-connection if and only if the following conditions are satisfied:*

$$(Cr1) \quad x \in M(\Gamma(y)) \Leftrightarrow y \in M(\Delta(x)), \quad x \in I, y \in A,$$

$$(Cr2) \quad \text{if } x \in M(\Gamma(y)), \text{ then the pair}$$

$$(\varepsilon_I(\Gamma(y), x), \varepsilon_A(\Delta(x), y))$$

is compatible with (Γ, Δ) .

Remark that the fundamental regular semigroup $U = U(I, A; \Gamma, \Delta)$ considered in [13] consists of all the pairs (f, g) that are compatible with (Γ, Δ) .

Recall from [16] that an isomorphism between two cross-connections $[I, A; \Gamma, \Delta]$ and $[I', A'; \Gamma', \Delta']$ is a pair of order isomorphisms $\alpha: I \rightarrow I'$, $\beta: A \rightarrow A'$, such that

$$\Delta = \alpha \circ \Delta' \circ \beta^\circ, \quad \Gamma = \beta \circ \Gamma' \circ \alpha^\circ,$$

where $\alpha^\circ: I'^\circ \rightarrow I^\circ$ and $\beta^\circ: A'^\circ \rightarrow A^\circ$ are defined as in Proposition 1. For any fundamental regular semigroup S , define $I_S = S/\mathcal{R}$, $A_S = S/\mathcal{L}$,

$$\lambda_e: I_S \rightarrow I_S, R_g \rightarrow R_{eg}, \quad e, g \in E(S),$$

$$\rho_e: A_S \rightarrow A_S, L_g \rightarrow L_{ge}, \quad e, g \in E(S),$$

$$\Gamma_S(L_e) = \ker \lambda_e, \quad \Delta_S(R_e) = \ker \rho_e, \quad e \in E(S).$$

Then $[I_S, A_S; \Gamma_S, \Delta_S]$ is a cross-connection, and S is isomorphic to a full regular subsemigroup of $U(I, A; \Gamma, \Delta)$ if and only if the cross-connection $[I_S, A_S; \Gamma_S, \Delta_S]$ induced by S is isomorphic to the cross-connection $[I, A; \Gamma, \Delta]$ (see [5, 16]).

An order preserving mapping $f: P \rightarrow Q$ of a poset P into a poset Q is said to be *residuated* if there exists an order preserving mapping $f^+: Q \rightarrow P$ such that $yf^+f \leq y$ and $x \geq xff^+$ for all $x \in P$ and $y \in Q$. The mapping f^+ is called the *residual* of f [2]. It is easy to see that if the residual f^+ of f exists, then it is unique. Further, if f and g are residuated mappings, and if fg exists, then it is a residuated mapping and its residual is g^+f^+ . It follows that the set $\text{Res } P$ of all residuated transformations of P is a semigroup under the composition of transformations, and that $f \rightarrow f^+$ is a dual isomorphism of

$\text{Res } P$ onto the semigroup $\text{Res}^+ P$ of all residuals of elements of $\text{Res } P$. It is clear that each $f^+ \in \text{Res}^+ P$ is a residuated transformation of P^{op} with residual $f \in \text{Res } P$. Thus there is a natural isomorphism between $\text{Res}^+ P$ and $\text{Res } P^{\text{op}}$ [8]. For convenience we shall henceforth identify $\text{Res}^+ P$ with $\text{Res } P^{\text{op}}$ and regard f^+ as a residuated transformation on P^{op} whenever $f \in \text{Res } P$. We shall also write elements of $\text{Res } P$ [$\text{Res } P^{\text{op}}$] as right [left] operators. It follows that $\text{Res } P \rightarrow \text{Res } P^{\text{op}}, f \rightarrow f^+$ is an isomorphism. For all $f \in \text{Res } P$ and $x \in P$ we thus have

$$(f^+ x)f \leq x \leq f^+(xf). \quad (3)$$

We say that $f \in \text{Res } P$ is *totally range closed* if f maps principal ideals onto principal ideals [2, 8]. Observe that a residuated transformation that is also normal must be totally range closed. Further, $f \in \text{Res } P$ is said to be *strongly range closed* if f and f^+ are totally range closed transformations of P and P^{op} respectively. It is clear that the set $B(P)$ of all strongly range closed transformations of P is a subsemigroup of $\text{Res } P$, and $f \rightarrow f^+$ is an isomorphism of $B(P)$ onto $B(P^{\text{op}})$. If $f \in \text{Res } P$, and if both f and f^+ are normal, then we shall say that f is *binormal*; if this is the case, then $f \in B(P)$. Again, the set of all binormal transformations of P forms a subsemigroup of $B(P)$.

PROPOSITION 3. *Let L be a complemented modular lattice, let $a \in L$ and let a' be a complement of a in L . Then*

$$(a; a'): L \rightarrow L, x \rightarrow (x \vee a) \wedge a' \quad (4)$$

is a binormal idempotent transformation such that

$$(a; a')^+ : L^{\text{op}} \rightarrow L^{\text{op}}, \quad y \rightarrow (y \wedge a') \vee a \quad (5)$$

is the residual of $(a; a')$. Further,

$$\begin{aligned} \ker(a; a') &= \Delta(a) = \{(x, y) \mid x \vee a = y \vee a\} \\ \ker(a; a')^+ &= \Gamma(a') = \{(x, y) \mid x \wedge a' = y \wedge a'\} \end{aligned} \quad (6)$$

and

$$M(\Gamma(a)) = M(\Delta(a)) = \{a' \mid a' \text{ is a complement of } a \text{ in } L\}. \quad (7)$$

Conversely, if e is any strongly range closed idempotent transformation of L , then e is binormal and $e = (a; a')$ for some $a \in L$ and some complement a' of a in L .

Proof. It is noted in [17] that $(a; a')$ is a normal transformation. We nevertheless provide a proof for the sake of completeness.

For any $x \in L$, we have

$$\begin{aligned}
 (a; a')^+(x(a; a')) &= (((x \vee a) \wedge a') \wedge a') \vee a \\
 &= ((x \vee a) \wedge a') \vee a \\
 &= (x \vee a) \wedge (a' \vee a) \quad (\text{by modularity}) \\
 &= x \vee a,
 \end{aligned}$$

and

$$((a; a')^+ x)(a; a') = x \wedge a'.$$

Since $(a; a')$ and $(a; a')^+$ are order preserving, it follows that $(a; a') \in \text{Res } L$ and $(a; a')^+$ is the residual of $(a; a')$.

To prove that $\phi = (a; a')$ is normal, consider $x \in L$ and let $z = (a \wedge x)' \wedge x$, where $(a \wedge x)'$ is a complement of $a \wedge x$. Then $z \vee a \geq z \vee (x \wedge a) = x \wedge ((x \wedge a)' \vee (a \wedge x)) = x$ by modularity. So $z \vee a \geq x \vee a$. Clearly $z \vee a \leq x \vee a$, whence $x \vee a = z \vee a$. Consequently $x\phi = z\phi$. Suppose now that $z_1, z_2 \in L(z)$. Then

$$\begin{aligned}
 z_1\phi \leq z_2\phi &\Rightarrow z_1\phi \vee a \leq z_2\phi \vee a \\
 &\Rightarrow z_1 \vee a \leq z_2 \vee a \quad (\text{by modularity}) \\
 &\Rightarrow (z_1 \vee a) \wedge z \leq (z_2 \vee a) \wedge z \\
 &\Rightarrow z_1 \leq z_2 \quad (\text{by modularity and since } a \wedge z = 0).
 \end{aligned}$$

It follows that ϕ is an order-isomorphism of $L(z)$ onto $L(z\phi) = L(x\phi)$. Thus $(a; a')$ is normal. By duality it follows that $(a; a')^+$ is a normal transformation of L^{op} . Thus $(a; a')$ is binormal. Also

$$(z_1, z_2) \in \ker(a; a') \Leftrightarrow z_1 \vee a = z_2 \vee a$$

as before, and so $\ker(a; a') = \Delta(a)$ is given by (6). Dually, $\ker(a; a')^+ = \Gamma(a')$ is given by (6).

Since for every complement a' of a , $(a; a')$ is a normal idempotent with $\ker(a; a') = \Delta(a)$ and $\text{im}(a; a') = L(a')$, it follows that every complement of a belongs to $M(\Delta(a))$. If $t \in M(\Delta(a))$, then $\phi = (a; a')$ is an isomorphism of $L(t)$ onto $L(a')$. In particular $(t \vee a) \wedge a' = t\phi = a'$, and so $a' \leq t \vee a$. Thus $1 = a \vee a' \leq t \vee a \leq 1$. Further $t \wedge a \in L(t)$ and $(t \wedge a)\phi \leq a\phi = 0$. Hence $0 = (t \wedge a)\phi$. Since $0, t \wedge a \in L(t)$ and since ϕ is an isomorphism when restricted to $L(t)$, we conclude that $t \wedge a = 0$. Thus t is a complement of a , and (7) holds.

If e is a strongly range closed idempotent transformation of L , then it follows from Theorem 13.4 of [2] that $e = (e^+0; 1e)$, and so by the above proof e is binormal.

Our result above shows that if L is a complemented modular lattice, then for every $a \in L$ there exist binormal idempotent transformations e and f such that $1e = a = f^+0$. The following shows that this property characterizes complemented modular lattices.

THEOREM 4. *For a poset I with 0 and 1 the following are equivalent:*

- (i) *I is a complemented modular lattice,*
- (ii) *for each $a \in I$ there exist binormal idempotent transformations e and f such that $1e = a = f^+0$,*
- (iii) *$B(I)$ is a regular semigroup, and*

$$\begin{aligned} B(I)/\mathcal{L} &\rightarrow I, & L_g &\rightarrow 1g \\ B(I^{\text{op}})/\mathcal{R} &\rightarrow I, & R_h &\rightarrow h0 \end{aligned}$$

are isomorphisms.

Proof. (i) \Rightarrow (ii) follows from Proposition 3.

(ii) \Rightarrow (iii). From Theorem 13.2 of [2] follows that I is a bounded lattice. From Theorem 14.5 of [2] and the corollary preceding it then follows that $B(I)$ is a regular semigroup. In view of (ii) we may conclude that $B(I)$ coordinatizes $B(I)$.

(iii) \Rightarrow (i) follows from Theorem 14.6 of [2].

Let L be a lattice with 0 and 1. For each $a \in L$ the relation

$$\Gamma(a) = \{(x, y) \mid x \wedge a = y \wedge a\} \quad (8)$$

is an equivalence relation on L , and the mapping

$$\Gamma: L \rightarrow \text{Eq}(L), \quad a \rightarrow \Gamma(a),$$

is an order preserving embedding of L into the poset $\text{Eq}(L)$ of all equivalence relations on L ordered under the reverse of inclusion. Note that $\Gamma(a) = \ker e_a$, where $e_a: L \rightarrow L$, $x \rightarrow x \wedge a$ is a normal retraction of L . Hence $\Gamma(a) \in L^\circ$ for all $a \in L$, and

$$\Gamma: L \rightarrow L^\circ, \quad a \rightarrow \Gamma(a) \quad (9)$$

is an order preserving embedding of L into L° . Proposition 3 shows that if L is a complemented modular lattice, Γ is an order preserving embedding of L into $(L^{\text{op}})^\circ$ also. Dually,

$$\Delta(a) = \{(x, y) \mid x \vee a = y \vee a\} \quad (10)$$

is a normal equivalence on L^{op} , and

$$\Delta: L^{\text{op}} \rightarrow (L^{\text{op}})^{\circ}, \quad a \rightarrow \Delta(a) \quad (11)$$

is an order preserving embedding of L^{op} into $(L^{\text{op}})^{\circ}$. Again by Proposition 3 we see that if L is a complemented modular lattice, then Δ is also an order preserving embedding of L^{op} into L° .

It may be noted that if L , Γ and Δ are given by the above, then $[L, L; \Gamma, \Gamma]$ and $[L^{\text{op}}, L^{\text{op}}; \Delta, \Delta]$ are the cross-connections induced by the semilattices L and L^{op} , respectively.

PROPOSITION 5. *Let L be a complemented modular lattice and let Γ and Δ be given by (8), (9), (10) and (11). For an order preserving transformation $f: L \rightarrow L$ the following statements are equivalent:*

- (i) f is normal and $\ker f = \Delta(a)$ for some $a \in L$,
- (ii) f is strongly range closed,
- (iii) f is binormal,
- (iv) there exists a normal transformation $g: L^{\text{op}} \rightarrow L^{\text{op}}$ such that (g, f) is compatible with (Γ, Δ) .

Moreover, if f satisfies these conditions, then f^+ is the unique normal transformation of L^{op} such that (f^+, f) is compatible with (Γ, Δ) .

Proof. (i) \Rightarrow (ii). Since f is normal, its image is a principal ideal of L . If $1f = b$, then $\text{im } f = L(b)$. Thus, for all $y \in L$, there exists $x \in L$ such that $xf = y \wedge b$. Since $\ker f = \Delta(a)$ we have $xf = zf$ if and only if $x \vee a = z \vee a$. Thus $\bigvee \{z \in L \mid zf = y \wedge b\}$ exists, and is equal to $x \vee a$ for some $x \in L$ with $xf = y \wedge b$. For $y \in L^{\text{op}}$, define f^+y by

$$f^+y = \bigvee \{z \in L \mid zf = y \wedge b\}.$$

Then $f^+: L^{\text{op}} \rightarrow L^{\text{op}}$ is well-defined. Further, we have

$$(f^+y)f = y \wedge b, \quad f^+(xf) = x \vee a$$

for all $x, y \in L$. If $y \leq y'$, then there exist $x, x' \in L$ with $x \leq x'$ such that $xf = y \wedge b$, $x'f = y' \wedge b$ since f is a normal transformation of L . Thus $f^+y = x \vee a \leq x' \vee a = f^+y'$. Thus f^+ is order preserving, and so f is residuated, f^+ being its residual. Since f is normal, it is also totally range closed. Obviously $\text{im } f^+$ is the principal ideal of L^{op} which is generated by a . Hence by Corollary 2 to Theorem 13.7 of [2], f is strongly range closed.

(ii) \Rightarrow (iii). If f is strongly range closed, then by Theorem 14.5 and Corollary 2 to Theorem 13.7 of [2] there exists a strongly range closed transformation $g: L \rightarrow L$ such that $f = f g f$ and $g = g f g$. Hence $e = f g$ is a

strongly range closed idempotent transformation, and $e = (a'; a)$ with $a = 1g$ and a' a complement of a in L by Proposition 3. Now $\alpha = f|L(a)$ is an isomorphism of $L(a)$ onto $L(b) = \text{im } f$, and $f = e\alpha$. By Proposition 3, e is normal. Hence f is normal. Similarly f^+ is strongly range closed, and so f^+ is also normal. We conclude that f is binormal.

(iii) \Rightarrow (iv). We show that (f^+, f) is compatible with (Γ, Δ) , that is, for all $x \in L$,

$$\Gamma(xf) = (f^+)^{\circ}(\Gamma(x)) = (f^+)^{-1}(\Gamma(x)),$$

and

$$\Delta(f^+x) = (\Delta(x))f^{\circ} = \Delta(x)f^{-1}.$$

Suppose that $(u, v) \in \Gamma(xf)$. Then $u \wedge xf = v \wedge xf$. Now f^+ preserves the meet operation (see [2, Exercise 4.2], or [8]). Hence $f^+(u \wedge xf) = f^+u \wedge f^+(xf)$. By Theorem 13.1* of [2], $f^+(xf) = x \vee f^+0$. Hence

$$\begin{aligned} f^+(u \wedge xf) &= f^+u \wedge (x \vee f^+0) \\ &= (f^+u \wedge x) \vee f^+0 \end{aligned}$$

by modularity. Now again by modularity

$$\begin{aligned} f^+u \wedge x &= (f^+u \wedge x) \vee (f^+0 \wedge x) \\ &= ((f^+u \wedge x) \vee f^+0) \wedge x \\ &= f^+(u \wedge xf) \wedge x \end{aligned}$$

and similarly

$$f^+v \wedge x = f^+(v \wedge xf) \wedge x.$$

From $u \wedge xf = v \wedge xf$ then follows $f^+u \wedge x = f^+v \wedge x$. Hence $(f^+u, f^+v) \in \Gamma(x)$, and thus $(u, v) \in (f^+)^{-1}(\Gamma(x))$.

If $(u, v) \in (f^+)^{-1}(\Gamma(x))$, then $f^+u \wedge x = f^+v \wedge x$. The binormal transformation f is in particular strongly range closed, so by Theorem 13.6 of [2],

$$u \wedge xf = (f^+u \wedge x)f = (f^+v \wedge x)f = v \wedge xf,$$

that is, $(u, v) \in \Gamma(xf)$. We conclude $\Gamma(xf) = (f^+)^{-1}(\Gamma(x))$. The equality $\Delta(f^+x) = \Delta(x)f^{-1}$ is proved dually.

(iv) \Rightarrow (i). If $g: L^{\text{op}} \rightarrow L^{\text{op}}$ is a normal transformation such that (g, f) is compatible with (Γ, Δ) , then

$$\Gamma(1f) = g^{-1}(\Gamma(1)) = \ker g$$

since $\Gamma(1)$ is the equality relation. Hence g satisfies the dual of (i), and from the above proof for (i) \Rightarrow (ii) we now see that g is strongly range closed, and from the proofs for (ii) \Rightarrow (iii) \Rightarrow (iv) we have that g is binormal and that (g^+, g) is compatible with (Γ, Δ) . Hence for all $x \in L$,

$$\Gamma(xg^+) = g^\circ(\Gamma(x)) = \Gamma(xf).$$

Since Γ is injective, $xg^+ = xf$ for all $x \in L$, thus $f = g^+$. Since g is binormal, $f = g^+$ must be normal. Further,

$$\Delta(g0) = \Delta(0)f^{-1} = \ker f$$

since $\Delta(0)$ is the equality relation. Hence (i) holds.

THEOREM 6. *Let L be a lattice with 0 and 1, and define Γ and Δ by (8), (9), (10) and (11). Then the following are equivalent:*

- (i) L is a complemented modular lattice,
- (ii) Δ is an order embedding of L^{op} into L° ,
- (iii) Γ is an order embedding of L into $(L^{\text{op}})^\circ$,
- (iv) $[L^{\text{op}}, L; \Gamma, \Delta]$ is a cross-connection.

If these conditions are satisfied, then the fundamental regular semigroup $U = U(L^{\text{op}}, L; \Gamma, \Delta)$ is given by

$$U = \{(f^+, f) \mid f \in B(L)\}. \quad (12)$$

Proof. That (i) implies (ii) and (iii) follows from Proposition 3, and that (iv) implies (ii) and (iii) follows from the definition of a cross-connection. It will be sufficient to prove that (ii) \Rightarrow (i) and (i) \Rightarrow (iv) hold: the proof for (iii) \Rightarrow (i) will follow by duality.

(ii) \Rightarrow (i). Let $a \in L$, and let $a' \in M(\Delta(a))$. Let e be the projection along $\Delta(a)$ upon $L(a')$. Since $\text{im } e = L(a')$ and $\ker e = \Delta(a)$, we may define

$$e^+y = \bigvee \{x \in L \mid xe = y \wedge a'\}, \quad y \in L,$$

and then

$$(e^+y)e = y \wedge a', \quad e^+(xe) = x \vee a, \quad x, y \in L,$$

so that e is a residuated transformation and e^+ its residual. Consequently $\text{im } e^+ = L^{\text{op}}(a)$, and from Theorem 13.4 of [2] we then have that $a' = 1e$ and $a = e^+0$ are complements in L . Thus L is complemented. Obviously e is a totally range closed idempotent transformation of L for all $a \in L$, $a' \in M(\Delta(a))$. Therefore by Corollary 1 to Theorem 13.7 of [2], (b, a) is a dual modular pair for all $a, b \in L$. Thus L is a modular lattice.

(i) \Rightarrow (iv). By Proposition 3, $a' \in M(\Gamma(a))$ if and only if a' is a complement of a , and then $a \in M(\Delta(a'))$. Thus the condition (Cr1) of Theorem 2 holds. By Proposition 5, $((a; a')^+, (a; a'))$ is compatible with (Γ, Δ) , since $(a; a')$ is binormal. Thus (Cr2) also holds and $[L^{\text{op}}, L; \Gamma, \Delta]$ is a cross-connection.

$U = U(L^{\text{op}}, L; \Gamma, \Delta)$ is the semigroup of all the pairs of normal transformations (g, f) , with $g: L^{\text{op}} \rightarrow L^{\text{op}}, f: L \rightarrow L$, that are compatible with (Γ, Δ) . Using Proposition 5, we must have (12).

COROLLARY 7. *If the equivalent conditions of Theorem 6 are satisfied, then*

$$U(L^{\text{op}}, L; \Gamma, \Delta) \rightarrow B(L), \quad (f^+, f) \rightarrow f \quad (13)$$

is an isomorphism.

If L is a complemented modular lattice, let $P(L)$ denote the subsemigroup of $B(L)$ which is generated by the idempotent transformations $(a; a')$, where a and a' are complements in L . Then from [17] we have that $P(L)$ is the idempotent-generated part of $B(L)$ and $E(P(L)) = E(B(L))$ consists of the pairs $(a; a')$, a and a' complementary in L . The biordered set $E(L)$ of idempotents of $B(L)$ (and of $P(L)$) was constructed in [17] in terms of the complemented modular lattice L . Using Theorem C of [13] one can construct yet another isomorphic copy $T(E(L))$ of $B(L) \cong U(L^{\text{op}}, L; \Gamma, \Delta)$.

Recall from [2, 7, 8] that the right annihilator X^r of a subset X of a semigroup S with zero 0 is given by

$$X^r = \{y \in S \mid xy = 0 \text{ for all } x \in X\}.$$

We write x^r instead of $\{x\}^r$. The left annihilators X^l and x^l are defined dually. A regular semigroup S with 0 is called a *strongly regular Baer semigroup* if the set of all principal left ideals of S coincides with the set of left annihilators of the elements of S and if the set of all principal right ideals of S coincides with the set of right annihilators of the elements of S . In other words, there exist surjective mappings

$$\sigma_l: S \rightarrow S/\mathcal{L}, \quad \sigma_r: S \rightarrow S/\mathcal{R},$$

such that for all $x \in S$,

$$S(x\sigma_l) = x^l, \quad (x\sigma_r)S = x^r.$$

From [2] we know that if this is the case, then S/\mathcal{L} and S/\mathcal{R} are dually isomorphic complemented modular lattices. It is easy to verify that if S is a strongly regular Baer semigroup, then every \mathcal{R} -coextension of a full regular

subsemigroup of S is also a strongly regular Baer semigroup. Further, if L is a complemented modular lattice, then $B(L)$ is a strongly regular Baer semigroup [8].

THEOREM 8. *A regular semigroup S is a strongly regular Baer semigroup if and only if the cross-connection $[I_S, A_S; \Gamma_S, \Delta_S]$ which is induced by S is isomorphic to the cross-connection $[L^{\text{op}}, L; \Gamma, \Delta]$ for some complemented modular lattice L .*

Proof. Let S be a strongly regular Baer semigroup. Then $L = S/\mathcal{L}$ is a complemented modular lattice. If μ is the greatest idempotent-separating congruence relation on S , then $S/\mu = T$ is a fundamental strongly regular Baer semigroup, and by Theorem 3 of [17] the idempotent-generated part of T is isomorphic to $P(L)$, i.e., the idempotent-generated part of $B(L)$. Consequently the biordered set of idempotents of S is isomorphic to the biordered set $E(L)$ of idempotents of $B(L)$. By Theorems B and C of [13] and by Corollary 7 we then have

$$U(I_S, A_S; \Gamma_S, \Delta_S) \cong T(E(S)) \cong T(E(L)) \cong B(L) \cong U(L^{\text{op}}, L; \Gamma, \Delta),$$

and so by Theorem 3.2 of [16], the cross-connection $[I_S, A_S; \Gamma_S, \Delta_S]$ is isomorphic to the cross-connection $[L^{\text{op}}, L; \Gamma, \Delta]$.

Conversely, if $[I_S, A_S; \Gamma_S, \Delta_S]$ is isomorphic to $[L^{\text{op}}, L; \Gamma, \Delta]$, then by Theorem 3.2 of [16], S/μ is isomorphic to a full regular subsemigroup of $U(L^{\text{op}}, L; \Gamma, \Delta)$ which in its turn is isomorphic to the strongly regular Baer semigroup $B(L)$. It follows that S is also a strongly regular Baer semigroup.

Remark 9. In view of the isomorphism $U(I_S, A_S; \Gamma_S, \Delta_S) \cong T(E(S))$ for any regular semigroup S , we could have stated the above result equivalently in terms of biordered sets as follows. A regular semigroup S is a strongly regular Baer semigroup if and only if the biordered set $E(S)$ of idempotents of S is isomorphic to the biordered set $E(L)$ of idempotents of $B(L)$ for some complemented modular lattice L .

The following corollary to Theorems 6 and 8 is analogous to Theorem 3 of [17].

COROLLARY 10. *Let L be a complemented modular lattice. The semigroup S is a fundamental strongly regular Baer semigroup coordinatizing L if and only if S is isomorphic to a full regular subsemigroup of $B(L)$, and $B(L)$ is determined up to isomorphism by this property.*

Remark 11. Let L be a complemented modular lattice. It follows from Theorem 1 of [17] that $E(L) = E(B(L))$ is a semilattice if and only if L is uniquely complemented, that is, if and only if L is a Boolean algebra (see

also Theorem 18.12 of [2]). In this case the idempotent $(a; a')$ of $B(L)$ is the normal retraction $e_{a'}: L \rightarrow L$, $x \rightarrow x \wedge a'$ onto the principal ideal $L(a')$ of L . The biordered set $E(L)$ which is induced by $B(L)$ is then isomorphic to the biordered set induced by the meet semilattice (L, \wedge) . The strongly regular Baer semigroup $B(L)$ which is isomorphic to $T(E(L))$ is then an isomorphic copy of the Munn semigroup $T(L)$ of the meet semilattice (L, \wedge) . In fact, the mapping

$$B(L) \rightarrow T(L), \quad f \rightarrow f | L((f^+0)')$$

is an isomorphism, where $(f^+0)'$ denotes the unique complement of f^+0 in L . Dually, $B(L)$ is also isomorphic to the Munn semigroup $T^{\text{op}}(L)$ of the join semilattice (L, \vee) . In view of this and Theorem 3.2 of [16] we have that the cross-connections $[L, L; \Gamma, \Gamma]$, $[L^{\text{op}}, L; \Gamma, \Delta]$ and $[L^{\text{op}}, L^{\text{op}}; \Delta, \Delta]$ are isomorphic.

If $\phi_i: S \rightarrow T_i$, $i = 1, 2$, are two representations of a semigroup S , we say that ϕ_1 is equivalent to ϕ_2 if there exists an isomorphism $\phi: \text{im } \phi_1 \rightarrow \text{im } \phi_2$ such that the following diagram commutes:

$$\begin{array}{ccc} S & \xrightarrow{\phi_1} & \text{im } \phi_1 \\ & \searrow \phi_2 & \downarrow \phi \\ & & \text{im } \phi_2 \end{array}$$

(D3)

Let S be a strongly regular Baer semigroup, let L_l be the poset of principal left ideals of S , and let L_r be the poset of principal right ideals of S . From [2, 8] we know that L_l and L_r are dually isomorphic complemented modular lattices. Further, for each $x \in S$

$$\phi_x: L_l \rightarrow L_l, \quad Sex \rightarrow ((Sex)^r)^l = Sex$$

is a strongly range-closed transformation on L_l and

$$\phi: S \rightarrow B(L_l), \quad x \rightarrow \phi_x \tag{14}$$

is a representation of S [2, 8]. The homomorphism ϕ is called the *right Janowitz representation* of S . For each $x \in S$,

$$\psi_x: L_r \rightarrow L_r, \quad eS \rightarrow ((xeS)^r)^l = xeS$$

is a strongly range closed transformation of L_r and

$$\psi: S \rightarrow B^{\text{op}}(L_r), \quad x \rightarrow \psi_x \tag{15}$$

is a representation of S [2, 8]. The homomorphism ψ will be called the *left Janowitz representation* of S , and

$$(\psi, \phi): S \rightarrow B^{\text{op}}(L_r) \times (L_l), \quad x \rightarrow (\psi_x, \phi_x) \quad (16)$$

the (two-sided) *Janowitz representation* of S .

Let $I_S = S/\mathcal{R}$, $A_S = S/\mathcal{L}$, and for $x \in S$, consider the transformations

$$\lambda_x: I_S \rightarrow I_S, \quad R_e \rightarrow R_{xe},$$

$$\rho_x: A_S \rightarrow A_S, \quad L_e \rightarrow L_{ex}.$$

Then $\lambda_x \in S^{\text{op}}(I_S)$ and $\rho_x \in S(A_S)$ for all $x \in S$ [4], and

$$\lambda: S \rightarrow S^{\text{op}}(I_S), \quad x \rightarrow \lambda_x, \quad (17)$$

$$\rho: S \rightarrow S(A_S), \quad x \rightarrow \rho_x \quad (18)$$

are representations of S ([4, 6]). It was observed in [5] that

$$(\lambda, \rho): S \rightarrow S^{\text{op}}(I_S) \times S(A_S), \quad x \rightarrow (\lambda_x, \rho_x) \quad (19)$$

maps S onto a full regular subsemigroup of $U(I_S, A_S; \Gamma_S, \Delta_S)$. We call λ [ρ , (λ, ρ)] the *left* [*right, two-sided*] *Hall representation* of S .

THEOREM 12. *Let S be a strongly regular Baer semigroup. Then the representations ψ , ϕ , (ψ, ϕ) , λ , ρ and (λ, ρ) which are given by (14), (15), (16), (17), (18) and (19) are equivalent and*

$$\ker \lambda = \ker \rho = \ker(\lambda, \rho) = \ker \psi = \ker \phi = \ker(\psi, \phi) = \mu, \quad (20)$$

where μ is the greatest idempotent separating congruence on S .

Proof. From Theorem 8 we know that the cross-connection $[I_S, A_S; \Gamma_S, \Delta_S]$ is isomorphic to a cross-connection $[L^{\text{op}}, L; \Gamma, \Delta]$, where L is a complemented modular lattice, and where Γ and Δ are given by (9) and (11). Since Γ and Δ are injective, we see that Γ_S and Δ_S are injective. Hence, by Lemma 3.9 of [16] and its dual we then have $\ker \lambda = \ker \rho = \ker(\lambda, \rho)$, where $\ker(\lambda, \rho) = \mu$ follows from the results of [4, 6]. Thus λ , ρ , and (λ, ρ) are equivalent representations.

Obviously $\kappa: L_l \rightarrow A_S$, $Se \rightarrow L_e$ is an order isomorphism, and for all $x \in S$, $\kappa\rho_x\kappa^{-1} = \phi_x$. From this we have that $\text{im } \rho \rightarrow \text{im } \phi$, $\rho_x \rightarrow \phi_x$ is an isomorphism. Analogously, $\text{im } \lambda \rightarrow \text{im } \psi$, $\lambda_x \rightarrow \psi_x$ is an isomorphism. From this and the above we find $\ker \phi = \ker \rho = \mu = \ker \lambda = \ker \psi$. Whence also $\ker(\psi, \phi) = \ker \psi \cap \ker \phi = \mu$. We conclude that (20) holds, and that all the above considered representations are equivalent.

In the following L will always denote a complemented modular lattice. We shall now investigate the fundamental strongly regular Baer semigroup $B(L)$.

THEOREM 13. *Let $e = (a'; a)$ be any idempotent of $B(L)$. Then the semigroup $eB(L)e$ is isomorphic to $B(L(a))$.*

Proof. Consider

$$\theta: B(L(a)) \rightarrow B(L), \quad f \rightarrow ef. \quad (21)$$

Clearly ef is a strongly range closed transformation of L and $\text{im } ef \subseteq L(a)$. Since e induces the identity transformation on $L(a)$, we have $efe = ef$, and so $f\theta \in eB(L)e$. If $f\theta = g\theta$, then for all $x \in L(a)$

$$xf = (xe)f = x(ef) = x(f\theta) = x(g\theta) = x(eg) = (xe)g = xg,$$

so that θ is injective.

If $e' = (b'; b) \leq e$ in the natural partial order on the set of idempotents of $B(L)$, then $b \leq a$ and $a' \leq b'$ in L , and $b_1 = b' \wedge a$ is a complement of b in $L(a)$. Further,

$$a' \vee b_1 = a' \vee (b' \wedge a) = (a' \wedge a) \vee b' = b'$$

by modularity. For all $x \in L$,

$$\begin{aligned} (x \vee b') \wedge b &= (x \vee a' \vee b_1) \wedge a \wedge b \\ &= (((x \vee a') \wedge a) \vee b_1) \wedge b \end{aligned}$$

by modularity, and thus

$$xe' = xee'',$$

where $e'' = (b_1; b)$ is an idempotent in $B(L(a))$. Now if f is any element of $eB(L)e$, then f has a normal factorization which is of the form $f = e'\alpha$, where $e' = (b'; b) \leq e$ and where α is an isomorphism among principal ideals of L that are contained in $L(a)$. By the above argument $f = ee''\alpha$, where $e'' = (b_1; b)$ as before. Since $e''\alpha \in B(L(a))$, it follows that θ is surjective.

Clearly θ is a homomorphism. We have proved that θ is an isomorphism of $B(L(a))$ onto $eB(L)e$.

The group of units of $B(L)$ is exactly $\text{Aut } L$, i.e., the automorphism group of L . We therefore have the following.

COROLLARY 14. *The maximal subgroup H_e of $B(L)$ corresponding to the idempotent $(a'; a) = e$ is isomorphic to $\text{Aut } L(a)$ (and to $\text{Aut } L^{\text{op}}(a')$).*

Recall from [8] that the elements $a, b \in L$ are called algebraically

equivalent, that is $a \sim_a b$, if and only if $L(a)$ is isomorphic to $L(b)$. We have the following characterization of Green's relation \mathcal{D} on $B(L)$ in terms of the algebraic equivalence on L .

THEOREM 15. *Let $f_1, f_2 \in B(L)$, $1f_i = a_i$, $f_i^+ 0 = b_i$, $i = 1, 2$. Then the following are equivalent.*

- (i) $f_1 \mathcal{D} f_2$ in $B(L)$,
- (ii) $a_1 \sim_a a_2$ in L ,
- (iii) $b_1 \sim_a b_2$ in L^{op} .

Proof. Let a'_i be a complement of a_i in L , $i = 1, 2$. Then $e_i = (a'_i; a_i) \mathcal{L} f_i$ in $B(L)$, $i = 1, 2$. Thus $f_1 \mathcal{D} f_2$ in $B(L)$ if and only if $e_1 \mathcal{D} e_2$ in $B(L)$. If $e_1 \mathcal{D} e_2$, then $e_1 B(L) e_1$ is isomorphic to $e_2 B(L) e_2$. Using Theorem 13 we then have

$$B(L(a_1)) \cong e_1 B(L) e_1 \cong e_2 B(L) e_2 \cong B(L(a_2)).$$

Therefore $L(a_1) \cong L(a_2)$ and thus $a_1 \sim_a a_2$. Conversely, if $a_1 \sim_a a_2$, then $L(a_1) \cong L(a_2)$, so that $B(L(a_1)) \cong B(L(a_2))$, and again by Theorem 13, $e_1 B(L) e_1 \cong e_2 B(L) e_2$. From [14] and from the fact that $B(L) \cong T(E(L))$ now follows that $e_1 \mathcal{D} e_2$ in $B(L)$. We have shown that (i) and (ii) are equivalent. Since $f_1 \mathcal{D} f_2$ in $B(L)$ if and only if $f_1^+ \mathcal{D} f_2^+$ in $B(L^{\text{op}})$, the equivalence of (i) and (iii) follows for dual reasons.

Remark 16. If L is a complemented modular lattice and $a, b \in L$, then we say that a and b are *perspective* if a and b have a common complement, and in this case we use the notation $a \not\sim b$. The reflexive symmetric relation $\not\sim$ is called the *perspective relation*, and its transitive closure is an equivalence relation which is called the *projectivity relation*. If $a, b \in L$, then a and b are said to be *projective*, i.e., $a \sim_\not\sim b$, if they are related in the projectivity relation. The idempotents $e = (a'; a)$ and $f = (b'; b)$ of $B(L)$ are \mathcal{D} -related in $P(L)$ if and only if $a \sim_\not\sim b$ in L [17]. Since $P(L)$ is a subsemigroup of $B(L)$ we always have $\not\sim \subseteq \sim_\not\sim \subseteq \sim_a$. In case L is coordinatized by a continuous regular ring, then $\not\sim = \sim_\not\sim = \sim_a$ (see, e.g. [10]); in this case $f_1 \mathcal{D} f_2$ in $B(L)$ if and only if $1f_1 \not\sim 1f_2$. This property holds for some projective geometries as well (see Theorem 2.27 of [1]).

Let S be a regular semigroup. For $x \in S$, denote the set of inverses of x in S by $V(x)$. Let $\pi(S)$ be the relation on S

$$\pi(S) = \{(x, y) \in S \times S \mid V(x) = V(y)\}. \quad (22)$$

The equivalence relation $\pi(S)$ is in general not a congruence relation. Yet if $\pi(S)$ is a congruence relation, then it is the greatest congruence relation on S whose idempotent congruence classes form rectangular bands. This is, e.g.,

the case with orthodox semigroups and with V -regular semigroups [12]. In [3, 17, 18] regular semigroups S are considered for which $\pi(S)$ is the identity relation on S . It is easy to see that if $\pi(S)$ is the identity relation on S , then the same holds for every \mathcal{H} -coextension of any full regular subsemigroup of S .

THEOREM 17. *If S is a strongly regular Baer semigroup, then $\pi(S)$ is the identity relation on S .*

Proof. Let L be a complemented modular lattice and suppose that $a, b \in L$ have the same set of complements in L . Since L is relatively complemented, $a \wedge b$ has a complement c in $L(b)$. Let $(a \vee b)'$ be a complement of $a \vee b$ in L , and put $d = c \vee (a \vee b)'$. Then

$$\begin{aligned} a \vee d &= a \vee c \vee (a \vee b)' \\ &= a \vee (a \wedge b) \vee c \vee (a \vee b)' \\ &= a \vee b \vee (a \vee b)' = 1, \\ a \wedge d &\leq (a \vee b) \wedge d \\ &= (a \vee b) \wedge (c \vee (a \vee b)') \\ &= c \vee ((a \vee b) \wedge (a \vee b)') = c \leq b \\ &\quad \text{(by modularity, and since } c \leq a \vee b). \end{aligned}$$

Hence $a \wedge d \leq a \wedge b$, $a \wedge d \leq c$, and thus $a \wedge d \leq (a \wedge b) \wedge c = 0$. Consequently d is a complement of a , and so d is a complement of b also. Hence

$$0 = b \wedge d \geq b \wedge c = c$$

from which $c = 0$. This implies $a \wedge b = b$. By symmetry, $a = b$.

Let us now consider the semigroup $B(L)$. Distinct \mathcal{L} -classes of $B(L)$ are of the form $L_{(a';a)}$ and $L_{(b';b)}$, where a and b are distinct elements of L , and where a' and b' are complements of a and b , respectively. From the above we can find a complement a'' of a in L that is not a complement of b in L or we can find a complement b'' of b in L that is not a complement of a in L . In the first case $(a''; a) \in E(L_{(a';a)})$ but $R_{(a'';a)} \cap L_{(b';b)}$ contains no idempotent, and in the second case $(b''; b) \in E(L_{(b';b)})$ but $R_{(b'';b)} \cap L_{(a';a)}$ contains no idempotent. Similarly, if $R_{(a;a')}$ and $R_{(b;b')}$ are distinct \mathcal{R} -classes of $B(L)$, then there exists an idempotent $(a; a'') \in E(R_{(a;a')})$ such that $L_{(a;a'')} \cap R_{(b;b')}$ contains no idempotent, or there exists an idempotent $(b; b'') \in E(R_{(b;b')})$ such that $L_{(b;b'')} \cap R_{(a;a')}$ contains no idempotent. From Lemma 2.1 of [11] we conclude that $\pi(B(L))$ is the identity relation on $B(L)$.

Let S be any strongly regular Baer semigroup coordinatizing the

complemented modular lattice L . Then S is an \mathcal{H} -coextension of a full regular subsemigroup of $B(L)$, and consequently $\pi(S)$ is the identity relation on S .

Remark 18. It follows from the foregoing theorem that a strongly regular Baer semigroup S is orthodox if and only if it is an inverse semigroup, and this is the case if and only if the complemented modular lattice S/\mathcal{L} is a Boolean algebra (see also [15] and Remark 11). In particular the multiplicative semigroup of a regular ring R is orthodox if and only if R is an abelian regular ring (see also [15]).

EXAMPLE 19. Let V be a vector space over a division ring D , and let $L = \mathcal{P}(V)$ be the projective geometry which is associated with V . Then the semigroup of all semilinear endomorphisms $S(V)$ can be shown to be a regular semigroup containing the semigroup $LT(V)$ of all linear endomorphisms of V as a full regular subsemigroup [9]. Hence the cross-connection induced by $S(V)$ is isomorphic to $[L^{\text{op}}, L; \Gamma, \Delta]$. By Theorem 8, $S(V)$ is a strongly regular Baer semigroup. Let μ be the greatest idempotent-separating congruence on $S(V)$. Then $S(V)/\mu$ is isomorphic to a full subsemigroup to $B(L)$. It can be shown [9] that in $S(V)$

$$\mu = \{(f, g) \in S(V) \times S(V) \mid g = \alpha f \text{ for some } \alpha \in D\},$$

and that for each $f \in S(V)$ the unique element of $B(L)$ determined by f is the transformation

$$\bar{f}: L \rightarrow L, \quad w \rightarrow f(w).$$

Here $\mathfrak{P}: S(V) \rightarrow B(L)$, $f \rightarrow \bar{f}$ is a canonical homomorphism which induces μ on $S(V)$. In this case \mathfrak{P} is the right Hall representation (or the right Janowitz representation) of $S(V)$. The μ -classes containing idempotents are isomorphic to the group of non-zero elements of D under multiplication. A maximal subgroup of $S(V)$ corresponding to a projection $e(w'; w)$ of V onto $w \in L$ is isomorphic to the group of all semilinear automorphisms of the vector space w , and the subgroup of $B(L)$ corresponding to the idempotent $\mathfrak{P}(e(w'; w)) = (w'; w)$ is the projective group of the geometry $\mathcal{P}(w)$. By the fundamental theorem of projective geometry (see Theorem 2.26 of [1]), all the elements $g \in B(L)$ for which $1g$ is of dimension greater than one belong to the image of $S(V)$ under \mathfrak{P} (see [9]).

When L is a projective geometry, the elements of $B(L)$ are called projective transformations or projective maps. Thus projective maps on L are binormal maps on L . Since the semigroup $B(L)$ uniquely determines and is determined by the geometry L , it is possible to study geometrical properties of L in terms of algebraic properties of the semigroup $B(L)$ or of the semigroup $P(L)$. When L is Desarguan, L is also uniquely determined

by the semigroup $S(V)$ of all semilinear endomorphisms, and several results on the group of semilinear isomorphisms of V and the general linear group on V have a natural extension to $S(V)$.

REFERENCES

1. E. ARTIN, "Geometric Algebra," Interscience, New York, 1957.
2. T. S. BLYTH AND M. F. JANOWITZ, "Residuation Theory," Pergamon, Oxford, 1972.
3. M. R. BROWN, D. W. HARDY, AND R. J. PAINTER, An embedding theorem for weakly inverse semigroups, *Semigroup Forum* **2** (1971), 332–335.
4. P. A. GRILLET, The structure of regular semigroups. I. A representation, *Semigroup Forum* **8** (1974), 177–183.
5. P. A. GRILLET, The structure of regular semigroups. II. Cross-connections, *Semigroup Forum* **8** (1974), 254–259.
6. T. E. HALL, On regular semigroups, *J. Algebra* (1973), 1–24.
7. M. F. JANOWITZ, Baer semigroups, *Duke Math. J.* **32** (1963), 85–96.
8. M. F. JANOWITZ, A semigroup approach to lattices, *Canad. J. Math.* **8** (1966), 1212–1223.
9. E. KRISHNAN AND K. S. S. NAMBOORIPAD, Semigroup of semilinear transformations and semigroup of projective transformations, to be published.
10. F. MAEDA, "Kontinuierliche Geometrien," Springer-Verlag, Berlin, 1958.
11. J. MEAKIN AND K. S. S. NAMBOORIPAD, Coextensions of regular semigroups by rectangular bands, I. *Trans. Amer. Math. Soc.* **269** (1982), 197–224.
12. W. D. MUNN, Fundamental inverse semigroups, *Quart. J. Math. Oxford* **21** (1970), 157–170.
13. K. S. S. NAMBOORIPAD, Relation between biordered sets and cross-connections, *Semigroup Forum* **16** (1978), 67–81.
14. K. S. S. NAMBOORIPAD, Structure of regular semigroups, I, *Mem. Amer. Math. Soc.* **22**, No. 224 (1979).
15. K. S. S. NAMBOORIPAD AND F. J. PASTIJN, V -regular semigroups, *Proc. Royal Soc. Edinburgh A* **88** (1981), 275–291.
16. K. S. S. NAMBOORIPAD AND F. J. PASTIJN, Regular involution semigroups, to appear.
17. F. J. PASTIJN, Biordered sets and complemented modular lattices, *Semigroup Forum* **21** (1980), 205–220.
18. B. M. SCHEIN, Catholic semigroups, in "Proceedings, Conference in Honor of A. H. Clifford, Tulane University, New Orleans (1979)," pp. 207–214.